

CHANGING THE CONVERSATION ABOUT BEHAVIORAL HEALTH

MENTAL ILLNESS

44.7 MILLION

adults in US experience some form of mental illness each year¹

10.4 MILLION

adults experience a serious mental illness¹

SUBSTANCE USE DISORDER

20.2 MILLION

adults have a substance use disorder each year²

10.2 MILLION

adults with a substance use disorder have a co-occurring mental illness²

THE NEED FOR MEDICATION ADHERENCE

50%

Less than half of people with mental illness take their medications as prescribed³

77%

Likelihood of symptom recurrence after 1 year for those with mental illness⁴

90%

Likelihood of symptom recurrence after 2 years for those with mental illness⁴

\$193.2
BILLION

Lost earnings experienced each year by those with serious mental illness⁵



AcuRiseID™

Elevating Mindful Adherence

AcuRiseID™ is a urinalysis that helps physicians and clinicians objectively monitor medication adherence in patients with mental illness. It provides the clinical clarity needed to understand medication's role in helping to overcome the challenges of substance use disorder and mental illness.

30

AcuRiseID includes more than 30 drugs in its substance use disorder profile including amphetamines, benzodiazepines, opiates and many more

40

AcuRiseID can detect more than 40 medications within its mental health profile including those used to treat schizophrenia, ADHD, anxiety, bipolar disorder, depression and more

For more information, visit www.aegislabs.com



References: **1.** National Institute of Mental Health. Mental Illness. National Institutes of Health Website. <https://www.nimh.nih.gov/health/statistics/mental-illness.shtml>. Updated November 2017. Accessed September 2, 2018. **2.** Center for Behavioral Health Statistics and Quality.(2015). Behavioral health trends in the United States:Results from the 2014 National Survey on Drug Use and Health (HHS Publication No. SMA 15-4927, NSDUH Series H-50). Retrieved from <http://www.samhsa.gov/data/>. **3.** García S, Martínez-Cengotitabengoa M, López-Zurbano S, Zorrilla I, López P,Vieta E, González-Pinto A. Adherence to Antipsychotic Medication in Bipolar Disorder and Schizophrenic Patients: A Systematic Review. J Clin Psychopharmacol.2016 Aug;36(4):355-71. **4.** El-Mallakh P, Findlay J. Strategies to improve medication adherence in patients with schizophrenia: the role of support services. Neuropsychiatr Dis Treat. 2015 Apr 16;11:1077-90. **5.** Insel TR. Assessing the economic costs of serious mental illness. Am J Psychiatry. 2008 Jun;165(6):663-5.